

The protective effect of cold acclimation on the low temperature stress of the lotus (*Nelumbo nucifera*)

XIN LI¹, DI ZHANG^{2*}, JUN XU³, JUN JIANG³, HONGWEI JIANG³

¹Administrative Station of Suzhou Wetland Protection, Suzhou, P.R. China

²Department of Landscape Science and Engineering, School of Design, Shanghai Jiao Tong University, Shanghai, P.R. China

³Institute of Agricultural Science of Taihu Lake District, Suzhou, P.R. China

*Corresponding author: zhangdi2013@sjtu.edu.cn

The authors are fully responsible for both the content and the formal aspects of the electronic supplementary material. No editorial adjustments were made.

Electronic supplementary material (ESM)

Table S1. The list of primers for qRT-PCR of *N. nucifera* rhizomes

Gene	Forward primer	Reverse primer
<i>CAT1</i>	AAAAGATGGATTGATGCTCTATCTGACC	CCAAAGACTTGTCAGCCTGAGACC
<i>GPX</i>	CACGACTTCACTGTCAAGGATGC	GCTCCGTGTAGTTCGAATTGGTC
<i>APX</i>	CGCCTACTTGCTTCGCGTCT	CTTGTCGTGGCCACTCCTCA
<i>MSD</i>	GGTGCCCTTGTTGGGCTAT	GGAACTAAATTTGGCCCCTTGG
<i>CSD</i>	CACATTTCAACCCAAATAACATGACAC	AATTATTGCCTCTGCCACTCCATC
<i>CBF2</i>	CCTATCTTGAAATCTTCAACAAGTCG	TTGTTGTTTGGGTGGTGGAGATT
<i>CRP2</i>	TCGGATGGGTTGCTTCCTTC	CGAGCCAATCTGGGAAGTGC
<i>COR</i>	CCTGCCATGCGAGGTAGAGG	TGGTCCTTCGGATCCACCAT
<i>Actin</i>	GCCTTGCAATCCACATCTGTT	GTTCCCAGGCATTGCTGACA